Hydrodynamic diffusion and its breakdown near AdS₂ fixed points

Blaise Goutéraux

Center for Theoretical Physics, CNRS, Ecole Polytechnique, Institut Polytechnique de Paris, France

Thursday November 19, 2020

String Theory Seminar, Institute for Theoretical Physics, Utrecht University



- Based on work to appear (soon!) with Daniel Areán (Universidad Autónoma de Madrid), Richard A Davison (Heriot-Watt University, Edinburgh) and Kenta Suzuki (Ecole Polytechnique, Palaiseau, France) (applying this Fall).
- Follow-up talk to my previous seminar here in Utrecht: 'Slow relaxation and diffusion in holography'.
- My research is supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 758759).
- All references and text in magenta contain hyperlinks.

- Universal description of interacting systems in the long wavelength $x \gtrsim \ell_{\rm eq}$, late time regime $t \gtrsim \tau_{\rm eq}$ ('small gradients').
- Based on a truncation of the dynamics to the relaxation of a few conserved densities, following from the symmetries of the system.
 Eg in a fluid: conservation of translations, rotations, particle number and possibly boosts.
- Provides an effective description of many interesting systems that cannot be described perturbatively: liquid phase of water, electrons in ultra-pure Graphene, Quark-Gluon-Plasma, superfluids, etc.

Classical diffusive hydrodynamics



- In this talk, I will focus on diffusive hydrodynamics: relaxation of the gradient of a conserved density ρ.
- Examples: particle number (chemical potential), energy (temperature gradient), shear momentum (transverse velocity), etc.

Classical diffusive hydrodynamics

• Conservation equation (Fick's law)

$$\partial_t \rho + \nabla_i j^i = 0$$

Constitutive relation

$$j^{i} = -\frac{D}{\chi} \nabla^{i} \rho + O(\nabla^{2}, \partial_{t} \nabla), \quad \chi = \frac{\partial \rho}{\partial \mu}$$

Decompose linear perturbations in plane waves

$$\rho(t,x) = \rho_0 + \delta \rho e^{-i\omega t + ikx}$$

• Retarded Green's function

$${\cal G}^{\sf R}_{
ho
ho}(\omega,k)=rac{i\chi Dk^2}{\omega+iDk^2}$$

• In this low frequency, low k approximation, single, diffusive pole

$$\omega = -iDk^2 + O(k^4) \quad \Rightarrow \quad G_{
ho
ho}(t,x) \sim
abla^2 rac{e^{-x^2/(4Dt)}}{t^{d/2}}$$

Limits of applicability: causality

• Causality: upper bound on the diffusivity [HARTMAN ET AL'17]

$$D \lesssim v^2 \tau_{
m eq}$$

- v: 'effective lightcone velocity'.
- Emergent infrared Lorentz invariance: expect v = c_{ir}.
- More generally:
 - Lieb-Robinson velocity?
 - Butterfly velocity?
 - Fermi velocity?
- More precise definition of local equilibration scales $\tau_{\rm eq}$, $\ell_{\rm eq}$?



* There are other limitations on the applicability of hydrodynamics (eg unstable frames, long time tails) but I won't discuss them in this talk.

Limits of applicability: convergence

• In principle, the hydro series can be pushed to any order in k

$$\omega = -i\sum_{n=1}^{+\infty}\omega_{2n}k^{2n}$$

Does this series converge?

- Hard question to answer in general. Use holography to compute the series: in real space (constitutive relation) [Heller&AL'13], [Heller&AL'20] and in Fourier space (dispersion relation) [WITHERS'18].
- In Fourier space, [WITHERS'18] showed that the radius of convergence k_{eq} of the series of the shear diffusive mode of the RN-AdS₄ black brane matches a singularity in the complex k plane.



Adapted from [Withers'18]

Limits of applicability: convergence

- Further recent confirmation that the convergence radius is set by the collision of the hydro mode with the nearest non-hydro mode by [JANSEN&PANTELIDOU'20].
- Further related studies and arguments for the above in [GROZDANOV&AL'19], [GROZDANOV&AL'19], [ABBASL&TAHERY'20].
- Are there cases where the convergence radius can be determined without referring to a specific microscopic theory?
- Yes, provided there is a hierarchy of scales: examples using holography.

Neutral translation-breaking black brane: model

• Classical solution to the action, [Bardoux&al'12], [Andrade&Withers'13]

$$S = \int d^4x \sqrt{-g} \left(R + 6 - rac{1}{2} \sum_{i=1}^2 \left(\partial \varphi_i \right)^2
ight)$$

Metric and matter fields (breaks translations homogeneously)

$$ds^{2} = -r^{2}f(r)dt^{2} + r^{2}dx_{i}^{2} + \frac{dr^{2}}{r^{2}f(r)}, \quad \varphi_{i} = mx^{i},$$

$$f(r) = 1 - \frac{m^{2}}{2r^{2}} - \left(1 - \frac{m^{2}}{2r_{0}^{2}}\right)\frac{r_{0}^{3}}{r^{3}} \quad \Rightarrow \quad T = \frac{3r_{0}}{4\pi}\left(1 - \frac{m^{2}}{6r_{0}^{2}}\right).$$

• At high temperature $T \gg m$, translations are weakly broken at a rate $\Gamma \sim m^2/T$ [DAVISON&GOUTÉRAUX'14]:

$$\partial_t P = -\Gamma P$$

Slow relaxation in the presence of weak explicit breaking

 Momentum couples to energy fluctuations: motion of poles governed by the equation

$$\omega^2 + i\Gamma\omega + c_s^2 k^2 = 0, \quad \Gamma \ll \Lambda_{uv}$$

in the scaling limit $\omega \sim k \sim \Gamma$.

• Crossover between diffusion of energy + weak relaxation of momentum when $k \lesssim k_{eq} \sim \Gamma$

$$\omega = -i\frac{c_s^2}{\Gamma}k^2 + \dots, \quad \omega = -i\Gamma + i\frac{c_s^2}{\Gamma}k^2 + \dots$$

and propagating modes when $k\gtrsim k_{\rm eq}\sim \Gamma$

$$\omega = \pm c_s k + \dots$$



Slow relaxation in the presence of weak explicit breaking

• We can be more precise about k_{eq} . The solutions to

$$\omega^2 + i\Gamma\omega + c_s^2k^2 = 0\,,$$

are

$$\omega_{\pm} = -i\frac{\Gamma}{2} \pm \sqrt{k^2 c_s^2 - \frac{\Gamma^2}{4}}$$

- The first non-analyticity is then at $(\omega_{\text{coll}}, k_{\text{coll}}) \simeq (-i\Gamma/2, \Gamma/(2c_s))$ $\Rightarrow (\omega_{\text{eq}}, k_{\text{eq}}) \simeq (\Gamma/2, \Gamma/(2c_s)).$
- Differently from [WITHERS'18], the dispersion relation is very well approximated by truncating to the first non-trivial terms in Γ and $k \Rightarrow$ analytical determination of the convergence radius: consequence of the hierarchy of scales $\Gamma \ll \Lambda_{uv}$.
- Different than the usual hydro expansion in k, which would diverge as $k \to k_{\rm eq}^-$

Slow relaxation in the presence of weak explicit breaking

- Even though energy diffusion ω = −i(c_s²/Γ)k² + ... can be augmented to incorporate slowly-relaxing momentum ω = −iΓ + ... when Γ ≪ Λ_{uv}, it formally breaks down at (ω_{eq}, k_{eq}).
- The diffusivity is D_ε ≃ c_s²/Γ and is naturally expressed in terms of a velocity and a timescale, which are directly related to the motion of poles in the complex frequency plane.
- The diffusivity can also be written in terms of the local equilibration scales:

$$D\simeq rac{c_s^2}{\Gamma}\simeq rac{1}{2}rac{\omega_{
m eq}}{k_{
m eq}^2}$$

• Valid in the regime when $\Gamma \ll \Lambda_{uv}$.

$$D \simeq \frac{1}{2} \frac{\omega_{\rm eq}}{k_{\rm eq}^2} = \frac{1}{2} v_{\rm eq}^2 \tau_{\rm eq} \,, \qquad \tau_{\rm eq} = \frac{1}{\omega_{\rm eq}} \,, \qquad v_{\rm eq} = \frac{\omega_{\rm eq}}{k_{\rm eq}} \,,$$

- Thanks to the hierarchy of scales $\Gamma \ll \Lambda_{uv}$, we could obtain a simple relation between a diffusivity, the local equilibration timescales and a characteristic velocity.
- What about more generally, when no symmetry is weakly broken, but still in the presence of a hierarchy of scales? Here, low temperature.
- Relation between diffusivity and 'chaos parameters' $D \simeq v_B^2/\lambda_L$ [BLAKE'16], valid for low temperatures?
- We will investigate this question using solvable models of transport: holography, SYK, focusing on the diffusion of energy and transverse momentum.

Neutral translation-breaking black brane: low T spectrum

 At low temperatures m >> T, translations are strongly broken and the only hydrodynamic mode is that of diffusion of energy:

$$\omega = -iD_{\epsilon}k^2 + O(k^4), \quad D_{\epsilon}(T \to 0) = \frac{\sqrt{3}}{\sqrt{2}}\frac{1}{m}$$

- The horizon geometry becomes $AdS_2 \times R^2$ at T = 0 ($r_e = \sqrt{6}m$).
- At small temperatures T ≪ m, the emergent SL(2,R)×SL(2,R) symmetry fixes the form of the IR retarded Green's function

$$egin{split} \mathcal{G}_{IR} \propto \mathcal{T}^{2\Delta(k)-1} rac{\Gamma\left(rac{1}{2}-\Delta(k)
ight)\Gamma\left(\Delta(k)-rac{i\omega}{2\pi T}
ight)}{\Gamma\left(rac{1}{2}+\Delta(k)
ight)\Gamma\left(1-\Delta(k)-rac{i\omega}{2\pi T}
ight)}\,, \ \Delta(k) = rac{1}{2} + \sqrt{rac{9}{4}+2rac{k^2}{m^2}}\,, \end{split}$$

and generates an infinite tower of gapped, IR modes

$$\omega_n = -i2\pi T(n + \Delta(0)) + O(k^2), \qquad n = 0, 1, 2, \dots$$

Neutral translation-breaking black brane: low T spectrum



 Red line: analytical approximation to the location of the hydro pole in the scaling limit ω ∼ T ∼ k² ∼ ε, for T/m = 1/1000:

$$\omega(k) = -i\epsilon\sqrt{\frac{3}{2}}\frac{k^2}{m}\left(1+\epsilon\frac{k^2}{m^2}+\epsilon^2\left(\frac{4\pi T^2}{3m^2}+\frac{k^4}{m^4}\right)+\ldots\right).$$

• Crosses IR poles at $\omega \simeq \omega_n = -i2\pi T(n+2)$, $k^2 \simeq k_n^2 = i\omega_n/D_{\varepsilon}$

• $T \ll m \Rightarrow$ agreement way beyond $k \ll T$ (usual hydro expansion).

Neutral translation-breaking black brane: avoided crossings



Avoided crossings (with vanishingly small gaps as $T \rightarrow 0$ rather than pole collisions) as a function of real k. The red line is the analytical approximation around $\omega = \omega_n + \delta \omega$, $k^2 = k_n^2 + \delta(k^2)$

$$(\mathcal{D}_n\delta(k^2)-i\delta\omega)(1-i\tau_n\delta\omega)-i\lambda_n\delta\omega=0,$$

$$\mathcal{D}_n \to D_{\varepsilon}, \quad \tau_n \to \frac{9m}{16\sqrt{6}(2+n)\pi^2T^2}, \quad \lambda_n \to \sqrt{\frac{3}{2}(n(n+4)+3)\frac{\pi T}{m}},$$

Neutral translation-breaking black brane: complex collision



The collision occurs for complex values of k (see also [Withers'18], [GROZDANOV&AL'19], [GROZDANOV&AL'19], [ABBASI&TAHERY'20], [JANSEN&PANTELIDOU'20])

$$\phi_k \to \frac{2^4}{6^{3/4}} \left(\frac{\pi T}{m}\right)^{3/2}, \quad k_{eq}^2 \equiv |k|^2 \to \frac{\omega_{eq}}{D_{\varepsilon}} \left(1 - \frac{4\sqrt{6}\pi T}{3m} + \ldots\right),$$
$$\omega_{eq} \equiv |\omega| \to 4\pi T \left(1 + \frac{8\sqrt{6}\pi T}{9m} + \ldots\right).$$

17

Neutral translation-breaking black brane: complex collision



The analytical approximations work very well

$$\phi_k \to \frac{2^4}{6^{3/4}} \left(\frac{\pi T}{m}\right)^{3/2}, \quad k_{eq}^2 \equiv |k|^2 \to \frac{\omega_{eq}}{D_{\varepsilon}} \left(1 - \frac{4\sqrt{6}\pi T}{3m} + \dots\right),$$
$$\omega_{eq} \equiv |\omega| \to 4\pi T \left(1 + \frac{8\sqrt{6}\pi T}{9m} + \dots\right).$$

Neutral translation-breaking black brane: diffusivity



$$\omega(k) = -i\epsilon \sqrt{\frac{3}{2}} \frac{k^2}{m} \left(1 + \epsilon \frac{k^2}{m^2} + \epsilon^2 \left(\frac{4\pi T^2}{3m^2} + \frac{k^4}{m^4} \right) + \dots \right).$$

Agreement between the approximation and the numerics implies

$$D_arepsilon(extsf{T} o 0) = rac{\omega_{ extsf{eq}}}{k_{ extsf{eq}}^2}$$

Relates hydrodynamic data at $(\omega, k \ll T)$ to data that mark the edge of validity of hydrodynamics $(|\omega| \simeq \omega_{eq} \sim T, |k| \simeq k_{eq} \sim \sqrt{Tm} \gg T)$

Reissner-Nordström black brane



- We find exactly similar results for the Reissner-Nordström black brane, both for energy diffusion and shear momentum diffusion: not specific to energy diffusion. Instead, hierarchy of scales $T \ll \mu$.
- The irrelevant deformation is different for the two diffusive modes

$$\Delta_{\varepsilon}(k=0)=2\,,\quad \Delta_{\Pi}(k=0)=1\,.$$

• Temperature dependence of the collision phase consistent with

$$\phi_k \sim T^{\Delta - 1/2}$$

Sachdev-Ye-Kitaev chain



• The Sachdev-Ye-Kitaev model ([SACHDEV&YE'93], [KITAEV'15]) and its higher-dimensional generalizations [GU&AL'16] are another set of solvable models of strongly-coupled matter.

$$H = i^{q/2} \sum_{x=0}^{M-1} \left(\sum_{\substack{1 \le i_1 < \ldots < i_q \le N}} J_{i_1 \ldots i_q, x} \chi_{i_1, x} \cdots \chi_{i_q, x} \right. \\ \left. + \sum_{\substack{1 \le i_1 < \ldots < i_{q/2} \le N}} J'_{i_1 \ldots i_q/2 j_1 \ldots j_{q/2}, x} \chi_{i_1, x} \cdots \chi_{i_{q/2}, x} \chi_{j_1, x+1} \cdots \chi_{j_{q/2}, x+1} \right).$$

 In the limit of infinite coupling J, J' → +∞, emergent reparameterization invariance suggests duality to near-AdS₂ gravity.

Exact large q solution



- By allowing q → +∞, the model can be solved analytically for all coupling strengths [CHOI&AL'20].
- The pole spectrum at strong coupling v → 1 is very close to the holographic results. One difference is that collisions occur for real p at strong enough coupling v ≥ 0.65.

Diffusivity relation



- We have reproduced their results and extracted the local equilibration scales.
- In the limit of strong coupling $v \to 1$, the $q = +\infty$ SYK chain also verifies

$$D_{arepsilon} = rac{\omega_{
m eq}}{k_{
m eq}^2}$$

Diffusivity bound

• Our results are compatible with the upper bound formulated by [HARTMAN ET AL'17]

$$D \lesssim v_{
m eq}^2 au_{
m eq}$$

with
$$v_{\rm eq} \equiv \omega_{\rm eq}/k_{\rm eq}$$
.

• In particular, they are compatible with the emergence of an 'effective lightcone velocity' v_{eq} even for non-relativistic systems.



Relation to chaos exponents

- In the examples of energy diffusion we studied, the diffusive approximation to the location of the pole is very good including at the energy pole skipping point [GROZDANOV&AL'17], [BLAKE&AL'18], [BLAKE&AL'18].
- This is the origin of the chaos relation

$$D_{\varepsilon} = v_B^2 / \lambda_L$$



Summary and outlook



 In states with a near-AdS₂ infrared fixed point, the excellent applicability of diffusive hydrodynamics across avoided crossings with an infinite tower of gapped infrared poles results in the relation

$$D = rac{\omega_{
m eq}}{k_{
m eq}^2}$$

where $\omega_{\rm eq}$ and $k_{\rm eq}$ are determined by infrared data, fixed by the symmetries of the state.

 As for the slow momentum-relaxing case, consequence of a hierarchy of scales.

Summary and outlook

- Extension to other near-AdS₂ states with non-universal leading irrelevant deformation [BLAKE&DONOS'16]?
- Addition of charge to the neutral, translation-breaking black brane: extra diffusive mode, governs the resistivity, of direct interest for strange metallic transport [HARTNOLL'14].
- Other types of fixed points with different scaling symmetries (Lifshitz, hyperscaling violation)?
- Other hierarchy of scales (eg angular momentum, magnetic field)?